

Prospects for biological control of field horsetail *Equisetum arvense* L. in New Zealand

Quentin Paynter, Jane Barton

Landcare Research
Private Bag 92170
Auckland
New Zealand

Landcare Research Contract Report: LC0708/100

PREPARED FOR: Horizons Regional Council

DATE: 2008-04-10

Reviewed by:

Approved for release by:

Lynley Hayes
Programme Leader
Landcare Research

Matt McGlone
Science Leader
Landcare Research

Stanley Bellgard
Scientist
Landcare Research

© Landcare Research New Zealand Ltd 2008

This information may be copied and distributed to others without limitation provided Landcare Research Limited and the source of the information are acknowledged. Under no circumstances may a charge be made to this information without the written permission of Landcare Research Limited.

Disclaimer

The findings in this report are specific to this project. Landcare Research accepts no responsibility where information in the report is used for any other purpose, and will not be liable for any loss or damage suffered as a result of such other use.

Contents

	Summary.....	4
1.	Introduction	6
2.	Background.....	6
	Global distribution and biology of field horsetail	6
	Distribution and status of field horsetail in New Zealand	8
	Current control methods	8
	Potential advantages and disadvantages of biological control	9
	Predicting establishment of biocontrol agents.....	9
	Predicting impact of biocontrol agents.....	10
3.	Objectives	11
4.	Methods	12
	Selecting a test-plant list for host-range testing of biological control agents.....	12
	Identifying fungal pathogens of field horsetail.....	12
	Identifying arthropod biocontrol agents of field horsetail.....	13
5.	Results	13
	The taxonomy of field horsetail.....	13
	Fungi.....	15
	Nematodes	17
	Arthropods.....	17
6.	Conclusions and recommendations	19
	Prospects for achieving biological control of <i>E. arvensis</i> in New Zealand.....	19
7.	Acknowledgements	20
8.	References	20
	Appendices	26

Summary

Project and Client

- The prospects for biological control of field horsetail *Equisetum arvense* in New Zealand were investigated by Landcare Research and Jane Barton, on behalf of Landcare Research, for Horizons Regional Council, funded by an Envirolink Medium Advice Grant (HZLC49).

Objectives

- Review the literature to identify potential biological control agents for field horsetail and to assess the feasibility of their release in New Zealand.
- Assess the prospects of achieving successful biocontrol of field horsetail in New Zealand.
- Estimate the total cost of the programme for the release of the agents.

Results

- No horsetail (*Equisetum*) species are native to New Zealand.
- Recent molecular analyses place the horsetails within, but only distantly related to, other ferns (Pteridophyta), having diverged by the end of the Devonian (~354 million years ago). The most closely related New Zealand native plants to *Equisetum* are believed to be the Marattoid ferns, of which there is one species, *Marattia salicina* (Para, King Fern). More distantly related native fern groups are the Ophioglossoid (5 species in two genera), Whisk (5 species in 2 genera), and Osmundaceous ferns (3 species in 2 genera).
- New Zealand Plantfinder (<http://www.plantfinder.co.nz/>, accessed 21/2/2008), listed no additional exotic Horsetail, Marattoid, Ophioglossoid, Whisk or Osmundaceous ferns that are sold commercially in New Zealand. All *Equisetum* species have been banned from sale now in New Zealand.
- Taxonomic isolation of *Equisetum* from native or valued exotic species in New Zealand indicates that relatively few test plant species should be required to demonstrate the host-range and environmental safety of candidate agents.
- Numerous arthropod and pathogens of field horsetail are present in the native range and many are apparently specific to *Equisetum*, such that there is a large pool of candidate agents likely to be suitable for use against field horsetail in New Zealand, including a flea beetle *Hippuriphila modeeri*, a weevil *Grypus equiseti*, numerous sawflies and several fungal plant pathogens, such as *Stammaria persoonii* and *Ascochyta equiseti*.
- Biological control programmes have not been conducted against field horsetail overseas, and little information exists on the impact of these natural enemies on field horsetail in its native range.

Conclusions and Recommendations

- A biocontrol programme is proposed based on the premise that the project should be approached from first principles. A survey should be undertaken to clarify what organisms are already present on field horsetail in New Zealand (estimated cost ~\$50,000–80,000, depending on how comprehensive). If nothing of significance is found, the control agents available in the native range of field horsetail should be assessed to develop a prioritised shortlist of agents for host-range testing. Several tasks have been identified that would narrow the search for effective control agents, and clarify the prospects for effective biological control of field horsetail in New Zealand:
 - Contact weed research organisations in other countries where field horsetail is invasive (e.g., Australia, South Africa) to investigate the possibility of collaboration on biological control of field horsetail.
 - Survey areas within the natural range of field horsetail that climatically most resemble New Zealand and where candidate agents that appear to have biocontrol potential are known to occur (e.g., Western Europe and the USA). The estimated cost of three 1-

week surveys (spring, summer and autumn) in Europe range from ~\$40,000 (for a limited survey in southern England; CABI Europe-UK) to \$80,000 (CSIRO European Laboratory, France). Both estimates assume samples would be sent to New Zealand for identification, which would be an additional cost.

- Ship the most promising arthropod agents into New Zealand quarantine for host-range testing &/or sub-contract an overseas agency (e.g., CSIRO or CABI) to conduct host-range testing on most promising pathogens (or send a pathologist overseas to test in USA, or Australia, or South Africa).
- The costs of host-range testing and preparing an application to ERMA for the release of a biocontrol agent (including ERMA fees that are currently \$30,000) are estimated at ~\$200,000–250,000 per agent (the cost of testing pathogens is likely to be greater than arthropods, because host-range testing would have to be conducted overseas). Note: this figure does not include the cost of mass-rearing and release of agents and is based on a purely operational programme (i.e. it does not include underpinning science, for example, to predict the impacts of biocontrol agents or to monitor agent establishment and impact, post-release). The cost of the programme will also be greater if molecular techniques are required to determine the origin of field horsetail in New Zealand or if it becomes necessary to plant out ‘trap plants’ in the native range, to ensure that populations of natural enemies that are capable of attacking field horsetail plants growing in New Zealand are identified. It is unusual for a single agent to provide sufficient control and it is likely that several would be required.

1. Introduction

Field horsetail *Equisetum arvense* L. is a vascular plant that reproduces by spores rather than seeds. The genus *Equisetum* includes ~15 species, commonly known as horsetails and scouring rushes, which are found throughout most of the world, but are not native to Australia and New Zealand (Webb et al. 1988). Traditionally, the horsetails and scouring rushes compose the entire class Equisetopsida, the sole member of the division Equisetophyta (Arthropophyta in older works), though a recent molecular analysis places the genus within the ferns (Pteridophyta), related to Marattiales (Pryer et al. 2004).

Three introduced *Equisetum* species became naturalised in New Zealand. According to Webb et al. (1995) two of these; *Equisetum fluviatile* L. and *Equisetum hyemale* L., were eliminated before they could become widely established, however, *E. hyemale* is still present, but rare in New Zealand (http://www.rnzih.org.nz/pages/nppa_075.pdf). In contrast, field horsetail, which is a Holarctic (i.e. native to temperate regions of Eurasia and North America) species, has become an aggressive weed in New Zealand, particularly where rainfall is moderate to high or in riparian sites.

Field horsetail develops extensive underground rhizomes which are resistant to herbicides so that, once established, it is extremely difficult and expensive to control (Webb et al. 1988). Biological control should, therefore, be a desirable control option. This report makes use of published literature reviews, internet web pages, and correspondence with botanical experts to provide information to allow decision-makers to decide if a biological control programme against field horsetail is feasible and should proceed, and to understand what such a programme would involve.

2. Background

Global distribution and biology of field horsetail

Field horsetail is native to Europe, Asia, and North America and has naturalised in Madagascar, South Africa, South America, Australia, and New Zealand (Large et al. 2006).

According to Doll (2001), field horsetail is a perennial with a spreading rhizome system that produces numerous shoots and tubers. The rhizomes are dark brown or blackish, 3–5 mm in diameter, covered with brownish hairs that give them a felt-like feel, and grow vertically to 1.8 m deep and horizontally to lengths of 25–50 cm. The horizontal rhizomes produce numerous shoots and form rounded tubers about 12 mm in diameter either singly or in pairs.

Plants produce two types of stems. Fertile (reproductive) stems (Fig. 1) appear in the early spring and are whitish to light brown, unbranched, hollow, c. 8 mm in diameter, cylindrical, leafless, jointed, and 15–30 cm long. The tips of fertile stems end in a yellowish to brownish spore-producing cone (called a strobilus), 12.5–31 mm long. Fertile stems wither and die once spores have been produced, usually by early summer.

Fig. 1. Sterile (left) and fertile (right) stems of *Equisetum arvense*

Image: The UniProt Consortium
The Universal Protein Resource (UniProt)
[Nucleic Acids Res. 36:D190-D195\(2008\).](https://pubmed.ncbi.nlm.nih.gov/16214821/)
<http://beta.uniprot.org/taxonomy/3258>



Sterile (vegetative) stems (Fig. 1) emerge later than the fertile stems and look like miniature pine trees with their plume-like branches. This appearance also explains the plant's common name: horsetail. Sterile stems are green, either erect or somewhat prostrate, 15–60 cm tall and composed of slender, grooved, hollow joints, 1–1.5 mm in diameter.

Field horsetail thrives in many habitats, and is as at home in wet, poorly drained areas of fields and grasslands, wet meadows, streams and other sites with high water tables as it is in well-drained sites in farm fields, orchards and nursery crops, and sites with sandy or gravel soil such as along roadsides, railway tracks and beaches. In general, field horsetail appears most commonly in acidic and wet soil conditions.

In its native range, field horsetail can become a weed of field and vegetable crops (e.g., Cloutier & Watson 1985). Movement of rhizomes or tubers on tillage implements is a common means of starting new populations. Simulations done to predict the rate of spread estimated that six years after introducing horsetail into an agricultural field, the weed will infest 2.5 acres (Cloutier & Watson, 1985).

Since the time of ancient Rome, field horsetail has been used in traditional medicine in Europe. Its main uses are as a diuretic (something that increases urine output), for osteoporosis (it contains silica deposits, which may be beneficial for bone thickening), and for wound healing (<http://www.nlm.nih.gov/medlineplus/>). Modern studies have shown that field horsetail produces compounds with antimicrobial activity, and that these compounds are active against a wide range of organisms including bacteria and fungi (Radulovic et al. 2005).

While the weed may be useful in small doses, in larger quantities it can be toxic (<http://www.nlm.nih.gov/medlineplus/>). The coarse stems are not often eaten by mammalian herbivores because of their silica deposits (the origin of the alternative name of “scouring rush” is due to the abrasiveness of these silica deposits). If they are eaten, acute thiamine deficiency can result in “equisetosis”. Horses are primarily affected, sheep to a lesser degree, and equisetosis is rarely fatal for cattle. Horses can be killed if large amounts of horsetail are consumed. Hay containing 20% or more horsetail produces symptoms in horses in 2–5 weeks. Symptoms include unthriftiness followed by weakness, “staggers”, nervousness, faulty vision, and difficulty in turning. In advanced stages, horses may “go down” and not be able to rise. Such animals are nervous and make frantic efforts to stand (Hill & Foland 1986). In late stages, muscular rigidity and constipation may be observed. In fatal cases, death is preceded by quiescence and coma.

Distribution and status of field horsetail in New Zealand

Field horsetail was first recorded in New Zealand in 1922 (Webb et al. 1988). Healy (1966) stated that field horsetail stem fragments were unintentionally imported into New Zealand as a contaminant in the roots of *Iris* sp. imported from Japan, although neither Roy et al. (1998) nor Webb et al. (1988) identify the origin of New Zealand field horsetail populations. It may be that the material from Japan was detected subsequent to an earlier introduction of unknown origin.

Webb et al. (1988) described field horsetail as an aggressive weed that is becoming well established in some areas, particularly where rainfall is moderate to high or where it can grow in riparian sites. According to Roy et al. (1998), field horsetail is of limited distribution in New Zealand, but has the potential to become more widespread. Nevertheless, infestations have been recorded from Kawhia, Havelock North, New Plymouth, Wanganui, and Lower Rangitikei in the North Island and from Marlborough, Nelson (including a 200-ha infestation near Karamea), the West Coast, Christchurch, and Dunedin in the South Island. In New Zealand, field horsetail now forms pure stands in a wide range of damp habitats, preventing the seedlings of native species from establishing and in some areas blocking and altering watercourses, causing flooding.

In the Manawatu-Wanganui Region, field horsetail is commonly spread by rhizome and root fragments (fragments as small as 2–3 mm are viable) in gravel and sand used in landscaping and road works (Craig Davey, Horizons Regional Council, pers. comm.). Furthermore, random discoveries from areas where no building mix has been deposited (e.g., from the tops of some catchments) implies long-distance spore dispersal is also occurring. Indeed, the recent rate of field horsetail spread has been described as “phenomenal and unstoppable” and some very large infestations already occur on the Wanganui River system, with the potential for further spread considered to be “vast” (Craig Davey, Horizons Regional Council, pers. comm.).

Field horsetail is designated an Unwanted Organism in New Zealand and all *Equisetum* species are listed under the National Pest Plant Accord. It is therefore an offence under Sections 52 and 53 of the Biosecurity Act to knowingly propagate, distribute, spread, sell, offer for sale or display (in a place where organisms are offered for sale or exhibited) field horsetail in New Zealand (<http://www.biosecurity.govt.nz/pest-and-disease-response/pests-and-diseases-watchlist/field-horsetail>).

Current control methods

Field horsetail’s extensive and deep rhizome system means that tillage and cultivation alone only destroy the top growth and delay reestablishment. Weed scientists in Canada hand-weeded an area with horsetail 16 times during one summer. The following year these plots looked identical to the control plot (Cloutier & Watson 1985). Furthermore, the extensive underground rhizomes are resistant to herbicides, making field horsetail extremely difficult to control (Webb et al. 1988). Field horsetail suppression (and possibly eradication) will only result if the appropriate mix of practices is done as a sustained effort for several seasons.

The best means of controlling this weed while it infests only small areas are through quarantine and mechanical and cultural methods. In New Zealand, the recommended control methods

(http://www.weedbusters.co.nz/weed_info/detail.asp?WeedID=110, accessed 6 November 2007) are:

1. Dig out and incinerate all parts and contaminated soil.

2. Spray (summer): metsulfuron-methyl 600g/kg (5g/10L) or Tordon Brushkiller (25ml/10L). Add penetrant.

Overseas, it has been noted that cultural practices such as improved drainage and adequate lime and fertilization programmes will help suppress horsetail infestations. Applying nitrogen fertilizer to grass crops is helpful because horsetail responds minimally to nitrogen while grass crops respond quickly and significantly, gaining a competitive advantage over field horsetail because the weed is shade sensitive (Andersson & Ludegardh 1999).

Potential advantages and disadvantages of biological control

Biological control could offer many advantages over current control methods for the management of field horsetail, and may be the only realistic option for large infestations given the difficulty of controlling this weed by conventional means. Classical biological control, if successful, is more cost-effective than other forms of control as it offers continuous action and self-dispersal to areas that are not likely to be targeted by other control programmes.

Despite its advantages, biological control may not be a “silver bullet”; although recent analyses have indicated the success rate of weed biocontrol programmes has been greater than previously supposed (Hoffmann 1995; McFadyen 1998; Fowler et al. 2000; Briese 2000), complete successes, where biological control is so dramatic that no other control methods are required, only account for approximately one-third of all completed programs (McFadyen 1998). Furthermore, although biological control is often perceived as an environmentally benign alternative to chemicals, there have been recently reported cases of damage to non-target plants (e.g., Louda et al. 2003). Nevertheless, the risk of failure and the impacts of non-target attack are likely to be minor, compared with the potential benefits. For example, Paynter and Flanagan (2004) showed that, of the weed biocontrol programmes that did not deliver ‘complete’ control, most resulted in ‘substantial’ or ‘partial’ control (i.e. where biological control contributes to management, but other control methods are still required to achieve adequate control). Examples of ‘partial control’ in New Zealand (listed by Fowler et al. 2000) include alligator weed (where biocontrol of floating weed mats is successful, but agents do not attack terrestrial infestations) and ragwort (excellent control in many regions, but not in high rainfall areas such as the west coast of the South Island). ‘Failure’ (i.e. an inability to find or establish control organisms, or an absence of agent impact) is a rare outcome of weed biocontrol programmes (Paynter & Flanagan 2004).

Furthermore, data on host use of 112 insects, three fungi, and one mite indicated that virtually all risk of non-target attack is borne by native plant species closely related to the target weeds and that the risk to native flora can be judged reliably before introduction (Pemberton 2000). Similarly, a survey in New Zealand indicated that the overall reliability of host-range testing in past weed biocontrol programmes was high. Only two cases of non-target attack were recorded on native plant species: both examples were of plants very closely related to the target weeds, both examples were predictable from host-range testing and in both cases the impacts of non-target attack were minor (Paynter et al. 2004; 2008).

Moreover, a recent economic analysis indicated that the overall weed biocontrol effort provided a strongly positive return on investment, with benefits provided by successful programmes far outweighing the total costs: for every dollar invested in weed biocontrol in Australia, a benefit of \$23.10 is generated (Page & Lacey 2006)

Predicting establishment of biocontrol agents

Reliably predicting the likelihood of establishment and impacts of introduced arthropods and pathogens on plant populations has long been a goal of weed biological control programmes

and research has been conducted to determine the importance of both release size (e.g., Memmott et al. 1998; 2005) and climate matching (e.g., van Klinken et al. 2003) in relation to biological control agent success.

The best predictor of establishment success of new organisms is the number and size of releases. In New Zealand, weed biological control agent establishment rates are now very high, largely due to the increased effort put into multiplication, release and monitoring in the extensive technology transfer programme that Landcare Research operates with regional councils, Landcare groups, Landcorp, local farmers' groups, the Department of Conservation, and forestry companies (Fowler et al. 2000).

Furthermore, in a broad sense, climate matching should rarely be problematic because co-evolved weed biological control agents should, generally, be adapted to the same climatic conditions as those to which their host-plants are adapted. However, van Klinken et al. (2003) noted that climate became an issue when agents were collected from a restricted part of the range of a plant species that established over a wide range of climatic and ecological conditions (a similar scenario could also occur if the native distribution of a plant was restricted, compared with its introduced range). Problems may also arise if the target country has an unusual climate that the plant can tolerate but in which the agent struggles. Given the vast native range of field horsetail, it should be prudent to choose agents that are adapted to regions where the climate is most similar to New Zealand. Therefore, for both climatic and logistical reasons, the relatively mild climatic regions of Western Europe would appear to be a more promising area in which to conduct native range surveys than, for example, Scandinavia, Siberia or Canada. Climatic matching could be further refined by, for example, using climate matching software (e.g., CLIMEX), but this may still require a detailed understanding of agent biology to be able to climate-match in detail.

Predicting impact of biocontrol agents

Estimating the likely impact of an exotic herbivorous organism in a new environment is more challenging than predicting the likelihood of establishment. In addition to climatic matching, an organism will face a host of other factors, such as predation, parasitism and competition, that might affect its ability to thrive in a new environment, as well as the ability of its host plant to compensate for attack. Furthermore, information regarding the growth of field horsetail in New Zealand compared with the native range is lacking, as is information regarding the impact of pathogens and invertebrate herbivory on field horsetail in the native range. Note, however, that this is not unusual before beginning a weed biological control programme.

Denoth et al. (2002) found that the success of biological control against weeds increased with the number of agents released, although they argued that this result might be because of the likelihood that the right control species is released increases with the number of agents released (lottery model), rather than because of the cumulative impact of multiple natural enemies. Certainly, spectacular biocontrol successes have been achieved with only one agent (Denoth et al. 2002) and a challenge for biocontrol practitioners is to identify the agent(s) that are most likely to impact on weed populations.

Crawley (1989) and Charudattan (2005) reviewed, respectively, the use of insect and plant pathogenic fungi in weed biological control. Crawley (1989) showed that certain insect groups have proved more successful at reducing host plant abundance than others: for example, ~50% of releases of insects in the *Dactylopiidae* (cochineal insects), *Curculionidae* (weevils) and *Chrysomelidae* (leaf-beetles) that established resulted in marked or complete control of the target weed, whereas all the programmes that successfully established insects in the *Cerambycidae* (longhorn beetles; 17 examples), *Coreidae* (leaf-footed bugs; 9 examples)

and *Carposinidae* (fruitworm moths; 7 examples) failed to control their target plants (table 8, Crawley 1989). Charudattan (2005) noted that weeds with robust capacity for vegetative regeneration (such as field horsetail) are more difficult to control than those that lack this trait. Indeed, field horsetail produces shoots during the entire summer (Cloutier & Watson 1985), which implies that either a multivoltine (multiple generation) agent or a combination of agents, acting in succession, will be required to inflict sustained damage throughout the growing season. Furthermore, in its native range, field horsetail can be a weed of field and vegetable crops (e.g., Cloutier & Watson 1985). One might speculate, that reduced plant competition, or elimination of natural enemies by pesticide use might contribute to an absence of regulation of field horsetail in such habitats. If so, it follows that while biocontrol may be likely to succeed in a range of habitats, additional control methods may still be required should field horsetail become an invasive weed of cropping systems in New Zealand.

Weed biological control agents, especially rust fungi, can be so specialised that they show host specificity within a given plant species. Matching the target host's susceptibility with the candidate pathogen's virulence is of utmost importance for biocontrol success with some agents since, for example, host-pathogen interactions at the species and subspecies levels are often governed by single-gene differences in rusts (e.g., varietal specificity; Charudattan 2005). For example, efficacy of the skeletonweed rust fungus *Puccinia chondrillina* varies from low to high, depending on weed forms (Burdon et al. 1981). Eriophyid mites can show similar levels of specificity: at least one biotype of St John's wort *Hypericum perforatum* appears to be resistant to the eriophyid mite *Aculus hyperici* in Australia (Mayo & Roush 1997). For certain candidate agents, to facilitate searching for natural enemies from the correct biotype(s) it would be prudent to determine which subspecies and forms of a weed are present in the introduced range, thereby reducing the potential for host-plant resistance. This can involve the use of molecular techniques to determine the origin of a weed. Some studies have utilised a cheaper method of directly testing the susceptibility of a weed to candidate agent species by planting out 'trap plants' collected in the weeds' introduced range and monitoring damage to them in the native range.

Finally, Charudattan (2005) also concluded that the stakeholders' perceptions of the effectiveness of a biocontrol programme can be unpredictable, leading to conflicting views of 'success'. We believe the aims of any biocontrol programme against field horsetail should therefore be clearly defined from the outset, so that success or failure can be assessed objectively against a well-defined goal.

3. Objectives

- Review the taxonomy of field horsetail in relation to native plant species and economically important exotic plant species present in New Zealand, to develop a test-plant list for host-range testing of biological control agents and to identify any potential barriers to the use of a biological control programme against field horsetail in New Zealand.
- Review existing literature on the natural enemies of field horsetail.
- Summarise the literature and current information available from researchers worldwide on the potential for biological control of field horsetail.

4. Methods

Selecting a test-plant list for host-range testing of biological control agents

Host-specificity testing is used to discard potential weed biological control agents that are likely to cause significant undesirable non-target damage to either native or valued exotic plants. The simplest tests are extremely robust ‘no-choice’ tests where arthropods are confined on a particular test plant and either feed or die, or, for pathogens, inoculation is attempted to see if infection can result. No-choice tests define the fundamental host-range of a particular species (all the plant species it can survive/complete development on; van Klinken & Heard 2000). However, for many herbivorous insects this fundamental host range is broader than the realised host range under field conditions, so more complex and costly tests, such as field cage oviposition tests, may be required.

A centrifugal phylogenetic method (Wapshere 1974) has long been used to determine the host-range of a potential biological control agent by sequentially testing plant taxa most closely related to the target weed followed by increasingly distantly related taxa until the host-range has been circumscribed. This approach is supported by recent advances in molecular techniques: host-shifts in lineages of specialist phytophagous insects are strongly linked to the evolution of host-plant lineages, and in particular plant chemistry. Such insects show a strong phylogenetic conservatism of host associations (see Briese 1996; Briese & Walker 2002). This pattern of strong phylogenetic conservatism in diet suggests the non-target plants of greatest risk are those closely related to known hosts (Futuyma 2000), and this has been validated by recent reviews of non-target attack by insect (Pemberton 2000; Briese & Walker 2002; Louda et al. 2003; Paynter et al. 2004) and fungal (Barton (née Fröhlich) 2004) weed biological control agents.

We, therefore, consulted the scientific literature to determine the taxonomic position of field horsetail to identify the most closely related plant species present in New Zealand that should be used to assess the risk of non-target attack when conducting host-range tests. To do this we searched the Web of Science[®] database for recent publications regarding the phylogeny of the genus *Equisetum*. We then referred to the most recent checklist of native New Zealand plants (de Lange et al. 2006) to identify the New Zealand plant species that are most closely-related to field horsetail.

Identifying fungal pathogens of field horsetail

A table was compiled of the fungi that have been reported associated with field horsetail. Information for this was obtained by searching computer databases and Internet sites. The most useful sources for this table were the USDA Fungus-host database or FDSM (which includes most New Zealand plant disease records) at <http://nt.ars-grin.gov/fungaldatabases/fungushost/FungusHost.cfm> and the Fungal Records Database of Britain and Ireland (FRDBI) at <http://194.203.77.76/fieldmycology/FRDBI/FRDBI.asp>. The IMI fungal herbarium was also searched online (at <http://194.203.77.76/herbIMI/index.htm>), but it listed no organisms associated with field horsetail. We also undertook Google searches using the key words ‘*Equisetum arvense*’ and ‘pathogen’ or ‘fungi’ or ‘equiseti’ (the species name for many fungi found associated with *Equisetum* hosts). Once a list had been created from internet sources, further information about each fungus was sought in the published literature. The NZFUNGI database (at <http://nzfungi.landcareresearch.co.nz>) was used to determine which species were already present in New Zealand. Each fungus was then assessed to determine whether it is likely to be damaging to field horsetail and whether it is likely to be sufficiently specific for use as a classical biological control agent. Note that

viruses and bacteria are not included in this survey because such organisms would be more challenging, and more expensive, to use for biocontrol than fungi.

Identifying arthropod biocontrol agents of field horsetail

Identifying candidate arthropod biocontrol agents of field horsetail was more difficult than identifying pathogens because there is no equivalent of the FDSM or FRDBI databases for all arthropod herbivores. However, a list of lepidoptera recorded from field horsetail was obtained from the Natural History Museum's world listing (<http://www.nhm.ac.uk/research-curation/projects/hostplants/>). We also searched CAB Abstracts and Google Scholar for field horsetail and sub-searching, first with the keyword "invertebrate*", and then with the keyword "herbivore". The abstracts were then examined and relevant pests added to the list. Useful information was found in the *Handbooks for the Identification of British Insects* series (many of which contain appendices with host-records), which listed species associated with field horsetail in Britain and Ireland.

We referred to checklists of New Zealand fauna to determine whether any of the pest species recorded feeding on/infecting field horsetail already occur in New Zealand.

5. Results

The taxonomy of field horsetail

Equisetum is the sole surviving genus (with two subgenera *Equisetum* and *Hippochaete*) in the 'horsetails and scouring rushes'. The horsetails were traditionally regarded as class Equisetopsida, which were considered to be 'fern allies', rather than true ferns (as below):

- Kingdom: Plantae
 - Division Tracheophyta (vascular plants)
 - Class Lycopsidea, (*fern-allies*) the clubmosses and related plants
 - Class Equisetopsida, (*fern-allies*) the horsetails and scouring-rushes
 - Class Psilopsida, (*fern-allies*) the whisk ferns
 - Class Filices, the true ferns
 - Class Spermatopsida (sometimes as several different classes of seed-plants)

Recent molecular analyses, however, place the horsetails within the ferns (Pteridophyta), with the most closely related New Zealand native plants to *Equisetum* currently believed to be the Marattoid ferns (Pryer et al. 2004; Wikström & Pryer 2005; Schuettpelz et al. 2006). Uncertainty remains regarding the exact relationships among horsetail, marattoid, and leptosporangiate ferns (Fig. 2). However, the horsetails are undoubtedly only distantly related to other ferns: using molecular techniques, Pryer et al. (2004) estimated that they diverged by the end of the Devonian (~354 million years ago); an estimate supported by the presence of fossil relatives of horsetails dating back to the late Devonian.

No *Equisetum* species are native to New Zealand and none are commercially important ornamental plants or crops in New Zealand. Due to this taxonomic isolation from native or valued exotics in New Zealand a relatively short test plant list should be sufficient to demonstrate the host-range and, therefore, environmental safety of candidate biocontrol agents. We propose that a test plant list should include a native New Zealand representative from each genus of the following major fern groups:

- **Marattoid ferns:** There is one native species of Marattoid fern in New Zealand (de Lange et al. 2006): *Marattia salicina* (Para, King Fern). An historical record of another

Marattoid fern *Angiopteris aurata* in New Zealand is considered to be the result of a mislabelled specimen (Brownlie 1959).

- **Ophioglossoid ferns:** Two genera of Ophioglossoid ferns are present in New Zealand (Allen 1982): *Botrychium*, represented by *Botrychium lunaria* (Moonwort), *B. biforme* and *B. australe* (Parsley fern) and *Ophioglossum*, represented by *Ophioglossum coriaceum* and *O. petiolatum* (de Lange et al. 2006).
- **Whisk ferns:** Two genera of whisk ferns are present in New Zealand (de Lange et al. 2006); *Tmesipteris*, represented by *Tmesipteris elongata*, *T. lanceolata*, *T. sigmatifolia*, and *T. tannensis* and *Psilotum*, represented by *Psilotum nudum*.
- **Osmundaceous ferns:** One species of *Todea* (*T. barbara*) and two species of *Leptopteris* (*L. hymenophylloides* and *L. superba*) are native to New Zealand (de Lange et al. 2006).

We also searched New Zealand Plantfinder (<http://www.plantfinder.co.nz/>, accessed 21/2/2008), which indicated that no additional exotic Marattoid, Ophioglossoid, Whisk or Osmundaceous ferns are sold commercially in New Zealand.




Figure 2. Fern phylogeny resulting from Bayesian analysis of combined 5-gene dataset (reprinted with authors' permission from Schuettpelz *et al.* 2006). F = ferns; L = leptosporangiate ferns; C = core leptosporangiates; major fern groups are also indicated with brackets at right. Filled numbered circles indicate nodes receiving both good posterior probability (≥ 0.95) and good maximum likelihood bootstrap (≥ 70) support; nodes with open circles did not receive good support from both measures.

Potential agents for the biological control of field horsetail in the New Zealand

Fungi

Classical biocontrol

Thirty-seven species of fungi were found associated with field horsetail (listed in Appendix 1). This substantial number probably reflects the fact that field horsetail has been used in traditional medicine for centuries and has, therefore, been relatively well-studied. Of these, 17 have no potential as classical biological control agents either because they are unlikely to be sufficiently damaging to be useful, or because they have a very broad host range, or both (see Appendix 1). There are 14 fungi listed in Appendix 1 for which we know too little to decide whether or not they could be useful for biocontrol. If any of these fungi were found during a pathogen survey of field horsetail, it would be worth testing their virulence and/or host range before rejecting them as possible agents. The remaining six fungi with biocontrol potential that are worthy of further discussion are: *Stamnaria persoonii* (Moug.) Fuckel; *Ascochyta equiseti* (Desm.) Grove; *Cylindrosporium equiseti* W.C. Liu et R.L. Bai; *Monodidymaria equiseti* (Dobrozhakova) U. Braun; *Heterosporium equiseti* H.C. Greene; and, *Mycosphaerella altera* (Pass.) House. These fungi are all either Ascomycetes (*Stamnaria* and *Mycosphaerella*) or the asexual stages (anamorphs) of Ascomycetes (the other four taxa). The Phylum Ascomycota is the largest group of fungi and includes saprobes, which live on dead organic material; parasites (esp. of plants), which obtain nutrients from living organisms; and, most of the fungi involved in forming lichens (a fungus/algae symbiosis). Most fungi can produce spores both sexually (through meiosis) and asexually (through mitosis). The characteristic that links all Ascomycetes is that when they produce spores sexually, they produce them in a microscopic sack called an ascus. The genus names *Ascochyta*, *Cylindrosporium*, *Monodidymaria* and *Heterosporium* are all applied to fungi reproducing asexually (anamorphs). However, there is good evidence linking these genera to the Ascomycetes, for example, some *Heterosporium* (*Cladosporium*) species also produce ascospores characteristic of Ascomycetes in the genus *Mycosphaerella* (Kirk et al. 2001). The four anamorph species of interest to us either do not produce ascospores on field horsetail, or they do, but no-one has yet linked their anamorph and teleomorph names (a common situation as such pairs usually look completely different from each other).

There have been more than 100 records of *Stamnaria persoonii* and its asexual state *Titaeospora equiseti* (Desm.) Vassiljevsky growing in association with *Equisetum* species (Cooper & Kirk 2006; Farr et al. 2008). While there are also two records of this fungus on unrelated hosts (*Ilex cornuta* and *Lagerstroemia indica*) in Florida (see Appendix 1), these records stand out, and it would be worth confirming their validity. While host range testing would be necessary before any fungus was considered for importation into New Zealand as a biocontrol agent, it would appear that *S. persoonii* is mostly, if not entirely, restricted to hosts in the genus *Equisetum* and would be sufficiently host-specific to be used safely here. Whether or not the fungus is capable of inflicting sufficient damage to field horsetail to be useful for its control is a matter that would require further investigation. Promisingly, it has been reported that *S. persoonii* caused a leaf spot and leaf blight on field horsetail in Korea, where the plant is valued for its medical properties (Shin 1994). Dr Shin reported that

infection was heavy, especially in wet places, and was sometimes associated with insect damage and/or infection with another fungus (a *Gloeosporium* species; Hyeon Dong Shin, Korea, pers. comm.). He said that affected plants “looked like they had been sprayed with herbicide”. Given that the damage reported by Dr Shin sounds quite obvious, it is surprising that most other records in the literature do not clearly state that the fungus is associated with disease symptoms. It is possible that the Korean fungus is a different strain to the European/North American one, and this would need to be investigated.

There are at least eight collections of this fungus stored in the Landcare Research Herbarium at Tamaki (PDD), three under the name *S. persoonii*; one under *T. equiseti*; two under *Titaospora detospora* (Sacc.) Bubák and two under *Cylindrosporium equiseti* (Desm.) Died (Shaun Pennycook, Landcare Research, pers. comm.). All were collected in Europe. Unfortunately, these collections are no longer viable, so this fungus could only be tested for New Zealand if a living culture/specimen could be found. No fungi in this genus have been used previously as classical biocontrol agents (Barton (née Fröhlich) 2004).

Ascochyta equiseti is the next most common fungus with biocontrol potential, with about 40 known records, all from *Equisetum* species. References consistently report that the organism is associated with disease symptoms (See Appendix 1). While no *Ascochyta* species have been used as classical biocontrol agents previously (Barton (née Fröhlich) 2004), the genus is biologically and morphologically very similar to the genus *Phoma*. An exotic strain of *Phoma clematidina* has been released in New Zealand for the biological control of old man’s beard *Clematis vitalba* but it does not appear to have persisted here and is believed it may have been out-competed by other pathogens already present in old man’s beard plant tissues. *Ascochyta* spores are generally small and moist and so tend to disperse small distances, unlike the dry, wind-dispersed spores of rusts. Therefore, if *A. equiseti* were to be used as a classical biocontrol agent in New Zealand, it might be necessary to make many releases, or even to apply inoculum as if it were a mycoherbicide, to aid its spread.

Cylindrosporium equiseti was only ‘discovered’ fairly recently (Lui & Bai 2001). Little information is available, but it is said to be associated with small leaf spots. The limited host range reported in the literature would need to be checked, as it may be an artefact of being little-studied.

There is also little information available about the last three fungi *Monodidymaria equiseti*, *Heterosporium equiseti* and *Mycosphaerella altera* (see Appendix 1). They have all been reported associated with disease symptoms, and only on *Equisetum* species, but with so few records in the literature, further testing would be vital. No fungi in the genera *Cylindrosporium*, *Monodidymaria*, *Heterosporium* or *Mycosphaerella* have been used previously as classical biocontrol agents (Barton (née Fröhlich) 2004). However, that does not mean they could not be useful. While most of the fungi used for classical biocontrol to date have been rusts (Basidiomycota, Uredinales), the remainder have mostly been, like all six of these fungi, Ascomycetes and their anamorphs. Rusts do make very good classical biocontrol agents, so it is unfortunate no rusts have been reported associated with field horsetail. Unfortunate, but not unexpected: rusts are usually associated with seed-plants, and (less frequently) true ferns (Bauer et al. 1999; Kirk et al. 2001). That said, a rust has been discovered associated with a *Selaginella* species (Bauer et al. 1999), and as the Selaginellaceae is an even more primitive plant group than the Equisetaceae (<http://plantphylogeny.landcareresearch.co.nz/>), it is possible there is a rust on field horsetail that has yet to be discovered.

To summarise, there are at least six pathogenic fungi that could be useful classical biocontrol agents for field horsetail in New Zealand, either alone or as part of a suite of agents. Four of these occur in the USA, three in Germany and the UK; France and Italy are each home to two

taxa. Thus, surveys for pathogens with biocontrol potential against field horsetail should be conducted in the US, and/or Western Europe.

Inundative biological control: developing a mycoherbicide

The development of a mycoherbicide (as outlined on the final page of this document) requires considerable field testing and other research. Consequently, this technique is usually limited to targets that cause a sufficiently large problem to justify the high developmental costs. Furthermore, mycoherbicides are normally developed using fungi that already occur in the country where they will be used. The exception to this rule is that sometimes a fungus introduced as a classical agent can have its dispersal and efficacy enhanced by being formulated and applied as a mycoherbicide *after* it has been introduced. For example, (Hennecke 2004) proposed that inoculum of the fungus *Phloeospora mimosae-pigrae* should be applied as a spray against the weed *Mimosa pigra*, after it was introduced to Australia, as unassisted it was struggling to keep up with the rapid growth of the weed.

For a mycoherbicide to be developed against field horsetail in New Zealand either a suitable fungus would already need to be here or one of the potential classical agents discussed above would need to be introduced and a decision made to enhance it. Of the 37 fungi listed in Appendix 1, ten already occur in New Zealand. However, none of these have potential as the active ingredient of a mycoherbicide because they are all either saprobes that would cause little damage to the weed (e.g., *Lachnella villosa* and *Lachnum virgineum*), or have very wide host ranges that would make it undesirable to broadcast their inoculum over large areas (e.g., *Verticillium dahliae*). While it has been argued that pathogens with broad host ranges can be used in mycoherbicides as long as the risks to non-target plants are understood (Bourdôt et al. 2003), this is more appropriate for weeds that are problematic in agricultural areas (where the diversity of other species is limited) than for environmental weeds such as field horsetail. A bioherbicide that could damage New Zealand native species would be of little use against field horsetail.

Nematodes

A plant pathogenic nematode, *Pratylenchus penetrans*, has been recorded from field horsetail. It is already present in New Zealand and has a huge host range, including the pasture grasses that compete with field horsetail, and is not, therefore, a suitable control agent for field horsetail.

Arthropods

We found records of 38 arthropod species that feed on field horsetail, of which some are clearly unsuitable for use as biological control agents on the basis of inadequate host-specificity (Appendix 2). However, according to literature host-records, 26 species are apparently sufficiently host-specific to be considered for use against field horsetail in New Zealand. We examine, below, whether this shortlist can be further refined, based on the likelihood that an agent will be sufficiently specific and damaging.

Mites: The eriophyid mite *Eriophyes equiseti* is likely to be a suitable biological control agent for field horsetail. Eriophyids have been considered to be ideal biological control agents due to the debilitating damage they cause to plants and their largely specialized feeding habits: over 80% of eriophyids found on weeds are monophagous (feed on a single host) (Rosenthal 1996). Two eriophyids have been used successfully as biological control agents against weeds (*Aceria chondrillae* against skeleton weed *Chondrilla juncea* in Australia and the USA and *A. malherbae* against field bindweed *Convolvulus arvensis* in the USA; Cullen et al. 1982; Rosenthal 1996). Another, *Aceria genistae*, was recently (February 2008) released in New Zealand for biological control of Scotch broom *Cytisus scoparius*. Although *Eriophyes equiseti* has only been recorded from Hungary (Farkas 1960), this perhaps indicates that few

acarologists have searched for this species (eriophyids are microscopic and unlikely to be identified by casual observation). We have not been able to ascertain the kind of damage caused by this mite but many eriophyids cause plant deformities, such as galls. The remaining candidate agents can be categorised according to the damage they cause (see below).

Sap-suckers: An aphid, two cicadellid leafhoppers and two delphacid planthoppers are apparently specific to *Equisetum* (Appendix 2). One aphid species, *Aphis chloris*, has been used as a weed biological control agent in Australia, but did not make an important contribution to St John's Wort *Hypericum perforatum* control (Briese & Jupp 1995). Aphids are the most important vectors of plant viruses (Burnett & Kawchuk 2002) and could conceivably transmit viral diseases to non-host plants during exploratory feeding and this risk of non-target damage might be expensive to research. Furthermore, aphids (or other sapsuckers that secrete honeydew) may attract ants which may then predate other insects present on the plant, potentially reducing the effectiveness of other agents (Thum et al. 1997). Therefore, careful consideration should be given before sap-suckers are used as weed biological control agents. Nevertheless, few cicadellids and delphacids are attended by ants (Delabie 2001), indicating they may be more suitable than the aphid *Macrosiphum equiseti* for biological control of field horsetail. A cicadellid *Zygina* sp. was introduced to control bridal creeper *Asparagus asparagoides* in Australia. In combination with an introduced rust fungus it caused major reductions of above- and below-ground biomass of the target weed (Morin & Edwards 2006).

Leaf/branch-miners: Several agromyzid flies in the genus *Lyriomyza* mine the narrow branches (Appendix 2) of field horsetail. Agromyzids are usually easy to rear and establish but they rank among the least successful of weed biocontrol agents: Crawley (1989) found no examples where they resulted in marked control of the target weed. The European agromyzid *Phytomyza vitalbae* has only a minor impact on old man's beard *Clematis vitalba* in New Zealand (Paynter et al. 2006), where it is parasitized by at least eight native and introduced parasitoids (Paynter et al. 2008). Agromyzids should therefore be given a low priority for biological control of field horsetail in New Zealand, unless surveys demonstrate they are unusually damaging or if their impact is likely to complement the action of another agent (for example, in combination with a stem-miner).

Stem and root-miners: The flea beetle *Hippuriphila modeeri* and weevil *Grypus equiseti* are both apparently specific to *Equisetum*; their larvae mining the main stems. *Hippuriphila modeeri* larvae "eat much pith, turning the outer surface of the stem black" (Robbins 1990). The moth *Loxoterma tiedemanniana* also mines field horsetail stems. However, given its distribution (Appendix 2), it could be a difficult and expensive to collect and it may not be well matched, climatically, to the areas where field horsetail is a weed in New Zealand. A review of successful and unsuccessful weed biocontrol agents indicated that the Chrysomelidae (Crawley 1989; see also Syrett et al. 1996) and weevils (Crawley 1989) have exceptionally good records of success in biocontrol programmes. Moreover, among the Chrysomelidae, various species of stem flea beetles have been extremely successful, for example: *Agasicles hygrophila* on alligator weed (Stewart et al. 2000); *Longitarsus jacobaeae* on ragwort *Senecio jacobaea* (McEvoy et al. 1991); *Apthona* spp. on leafy spurge *Euphorbia esula* (e.g., Kirby et al. 2000) – the latter two being stem- and root-miners as larvae. This success implies that both the horsetail flea beetle *Hippuriphila modeeri* and the horsetail weevil *Grypus equiseti* should be high priority species for consideration as biological control agents of field horsetail in New Zealand.

Defoliators: According to host-records, approximately 15 species of sawfly graze the fronds of, and are apparently specific to, *Equisetum* (Appendix 2). Some, such as *Dolerus bimaculatus* are restricted to relatively cold, northern localities and some, such as *D. pratensis* and *D. aericeps*, are rare or local in Britain (Benson 1952).

however, have been described, respectively, as abundant in Great Britain and very common in England (Benson 1952), indicating they must be significant defoliators in the native range. Some adult sawflies are carnivorous (Jervis & Vilhelmsen 2000); although the *Dolerus* species listed by Jervis & Vilhelmsen (2000) were said to feed on pollen, nectar, honeydew and the sugar-containing spermatial fluid of rust fungi, it would be prudent to confirm the adult feeding preferences of any *Dolerus* species being considered for introduction to control field horsetail.

Five exotic sawfly species in the superfamily Tenthredinoidea have been accidentally introduced into New Zealand and become abundant: *Pontania proxima* and *Nematus oligospilus*, which attack willows; *Caliroa cerasi*, which attacks pear, cherry and hawthorn trees; and the raspberry sawfly *Priophorus morio*; and an Australian species *Phylacteophaga froggatti* mines eucalyptus leaves. Therefore, sawfly defoliators should be very promising candidate agents for the biological control of field horsetail in New Zealand. However, one sawfly *Monophadnus spinolae* has been used for biocontrol of an exotic weed (old man's beard *Clematis vitalba*) in New Zealand and its establishment has not been confirmed, even though the first release of this agent was performed in 1996. Mass rearing of the sawfly was difficult due to a 20:1, male: female sex ratio, asynchronous emergence, and poor larval survival (Gourlay et al. 2000).

6. Conclusions and recommendations

Prospects for achieving biological control of field horsetail in New Zealand

We conclude that the prospects of finding sufficiently specific classical biological control agents for introduction into New Zealand are extremely good. However, as field horsetail has not been targeted for biological control overseas, a biological control programme would have to start from scratch. Nevertheless, field horsetail has the potential to be a relatively inexpensive target for biological control because:

- As field horsetail is taxonomically distant from New Zealand native plants, an extensive test-plant list is not required and relatively simple laboratory no-choice experiments to determine the fundamental host-range are likely show that the candidate biocontrol agent is sufficiently safe for release in New Zealand. This means costly (and sometimes logistically difficult, or even impossible) field host-range tests are unlikely to be necessary. For arthropods, these tests could be carried out in quarantine in New Zealand. For plant pathogens, host-range testing would have to be conducted overseas as there is currently no pathogen-proof quarantine facility in New Zealand.
- Extensive literature records indicate that a large number of apparently specific arthropods and plant pathogens have already been recorded attacking field horsetail, so extensive surveys in the native range should not be required. Nevertheless, some surveys in the native range (especially in Western Europe and/or the USA) would be prudent to determine which agents appear to be the most damaging and widespread and to investigate whether agents are likely to complement one another by attacking different plant parts and/or by attacking plants at different times of the year.
- Significant opposition to biological control is highly unlikely because, while field horsetail is used in homeopathic preparations (Roy et al. 1998), it is not grown for this purpose in New Zealand (indeed it would be illegal to do so) and it is not valued by other sectors in New Zealand (e.g., as a fodder crop or a nectar source for beekeeping). While horsetails are still grown as ornamentals in some gardens this is unlikely to present a major stumbling block since they can no-longer be sold or propagated.

Two potential complications could arise because field horsetail has a huge native range, over which two subspecies (*ssp. arvense* and *ssp. boreale*) and numerous named forms and varieties occur (<http://data.gbif.org/search/taxa/Equisetum%20arvense>): (1) ensuring a good climate match between New Zealand and the area in which to search for potential agents and; (2) the potential for host-plant resistance.

To ensure good climate matching, it should be useful to determine the areas within the natural range of field horsetail that, climatically, most resemble New Zealand. This could be achieved using CLIMEX software. However, given that most weed biological control agents have been successfully introduced into New Zealand (directly or indirectly) from Western Europe (Hayes 2005), where field horsetail is native, searching for agents in this region is likely to result in an adequate climate match.

As noted in section 2, above, weed biological control agents, especially rusts and eriophyid mites, can be so specialised that they show host specificity within a given plant species. However, no rusts have been recorded attacking field horsetail (Appendix 1); indeed many of the most promising pathogens and insects attack more than one *Equisetum* species (Appendices 1, 2). We therefore predict that host plant resistant genotypes are unlikely to be an issue except, perhaps, for the eriophyid mite *Eriophyes equiseti*.

Recommendations

- Survey the invertebrate fauna and pathogens of field horsetail in New Zealand. Estimated cost is \$50,000–80,000.
- Contact weed research organisations in other countries where field horsetail is invasive (e.g., Australia, South Africa) to investigate the possibility of collaboration on biological control of field horsetail.
- Survey field horsetail localities in Britain and/or France and Germany and/or the USA and prioritise a shortlist of agents for host-range testing. Approximate cost of three 1-week surveys (Spring, Summer and Autumn) \$40,000 (CABI Europe-UK)–\$80,000 (CSIRO European Laboratory). Both estimates assume samples would be sent to New Zealand for identification, which would be an additional cost.
- Ship most promising arthropod agents into New Zealand quarantine for host-range testing &/or sub-contract an overseas agency (e.g., CSIRO or CABI) to conduct host-range testing on most promising pathogens (or send a pathologist overseas to carry out testing in USA, or Australia, or South Africa).

7. Acknowledgements

We thank Shaun Pennycook for helpful assistance. Simon Fowler and Stan Bellgard refereed an earlier draft of this report. This report was funded by a Foundation for Research, Science & Technology (FRST) Envirolink Medium Advice Grant (HZLC49).

8. References

- Allen HH 1982. Flora of New Zealand. Volume 1. Wellington, Government Printer.
- Andersson TN, Ludegardh B 1999. Growth of field horsetail (*Equisetum arvense*) under low light and low nitrogen conditions. *Weed Science* 47: 41–46.
- Aucejo S, Foo M, Gimeno E, Gomez-Cadenas A, Monfort R, Obiol F, Prades E, Ramis M, Ripolles JL, Tirado V, Zaragoza L, Jacas JA, Martinez-Ferrer MT 2003. Management of *Tetranychus urticae* in citrus in Spain: acarofauna associated to weeds. *Bulletin OILB/SROP*, 26 (6): 213–220.

- Barton (née Fröhlich), J. 2004. How good are we at predicting the field host-range of fungal pathogens used for classical biological control of weeds? *Biological Control* 31 (1): 99–122.
- Bauer R, Vánky K, Begerow D, Oberwinkler F 1999. Ustilaginomycetes on Selaginella. *Mycologia* 91 (3): 475–484.
- Benson RB 1952. Hymenoptera. 2, Symphyta. Section (b): Family Tenthredinidae. Handbooks for the identification of British insects. Royal Entomological Society of London. Pp. 51–137.
- Bourdôt GW, Hurrell GA, de Jong MD 2003. Defining safety zones for bioherbicides based on plurivorous pathogens using models of spore escape and dispersion. Paper read at Biocontrol of weeds with pathogens. Workshop before 8th International Congress of Plant Pathology, 1 February 2003, Lincoln, Christchurch, New Zealand.
- Braun U 1994. Miscellaneous notes on phytopathogenic hyphomycetes. *Mycotaxon* 51: 37–68.
- Briese DT 1996. Phylogeny: can it help us to understand host choice by biological weed control agents? In: Moran VC, Hoffman JH eds Proceedings of the IX symposium on biological control of weeds. South Africa, University of Cape Town. Pp. 63–70.
- Briese DT 2000. Classical biological control. In: Sindel BM ed. Australian weed management systems. Meredith, Australia, RG & FJ Richardson.
- Briese DT, Jupp PW 1995. Establishment, spread and initial impact of *Aphis chloris* Koch (Hemiptera: Aphididae), introduced into Australia for the biological control of St John's Wort. *Biocontrol Science and Technology* 5: 271–286.
- Briese DT, Walker A 2002. A new perspective on the selection of test plants for evaluating the host-specificity of weed biological control agents: the case of *Deuterocampta quadrijuga*, a potential insect control agent of *Heliotropium amplexicaule*. *Biological Control* 25: 273–287.
- Brownlie G 1959. Some problems in New Zealand fern nomenclature. *Transactions of the Royal Society of New Zealand* 87: 195–198.
- Bullock, JA 1992. Host plants of British beetles: a list of recorded associations. Brentwood UK, Cravitz Publishing Company.
- Burdon JJ, Groves RH, Cullen JM 1981. The impact of biological control on the distribution and abundance of *Chondrilla juncea* in south-eastern Australia. *Journal of Applied Ecology* 8: 957–966.
- Burnet PA, Kawchuk LA 2002. Insect-vectored crop diseases. In: Pimentel D ed. Encyclopedia of pest management. New York/Basel, M. Dekker. Pp. 407–409.
- Çalmaşur Ö, Özbek H 2004. A contribution to the knowledge of Tenthredinida (Symphyta, Hymenoptera) fauna of Turkey Part II: subfamilies Blennocampinae, Dolerinae, Nematinae and Selandrinae. *Turkish Journal of Zoology* 28: 55–57.
- Camara MPS, Palm ME, van Berkum P, O'Neill NR 2002. Molecular phylogeny of *Leptosphaeria* and *Phaeosphaeria*. *Mycologia* 94: 630–640.
- Charudattan R 2005. Ecological, practical, and political inputs into selection of weed targets: What makes a good biological control target? *Biological Control* 35: 183–196.
- Cloutier D, Watson A 1985. Growth and regeneration of field horsetail (*Equisetum arvense*). *Weed Science* 33: 358–365.
- Cooper J, Kirk PM 2006. The fungal records database of Britain and Ireland (FRDBI). <http://194.203.77.76/fieldmycology/FRDBI/FRDBI.asp> [accessed 12 February 2008].
- Crawley MJ 1989. The successes and failures of weed biocontrol using insects. *Biocontrol News & Information* 10: 213–223.
- Cullen JM, Groves RH, Alex JF 1982. The influence of *Aceria chondrillae* on the growth and reproduction capacity of *Chondrilla juncea*. *Journal of Applied Ecology* 19: 529–537.
- de Lange PJ, Sawyer JWD, Rolfe JR 2006. New Zealand indigenous vascular plant checklist. Wellington, New Zealand Plant Conservation Network.
- Delabie JHC 2001. Trophobiosis between Formicidae and Hemiptera (Sternorrhyncha and Auchenorrhyncha): an overview. *Neotropical Entomology* 30: 501–516.

- Dennis RWG 1986. Fungi of the Hebrides. Kew, UK, Royal Botanic Gardens.
- Denoth M, Frid L, Myers JH 2002. Multiple agents in biological control: improving the odds? *Biological Control* 24: 20–30
- Doll J 2001. Biology and control of field horsetail (*Equisetum arvense* L., Horsetail Family). http://128.104.239.6/uw_weeds/extension/articles/conhorsetail.htm [accessed when?]
- Ellis MB 1971. *Dematiaceous Hyphomycetes*. Kew, UK, C.A.B., Commonwealth Mycological Institute.
- Farkas HK 1960. Über die Eriophyiden (Acarina) Ungarns I. *Acta Zoologica Academiae Scientiarum Hungaricae* 6: 315–339.
- Farr DF, Rossmann AY, Palm ME, McCray EB 2008. Fungal Databases, Systematic Mycology and Microbiology Laboratory, ARS, USDA. <http://nt.ars-grin.gov/fungaldbases/> 2008 [accessed DATE February 2008; cited February/March 2008].
- Fowler SV, Syrett P, Hill RL 2000. Success and safety in the biological control of environmental weeds in New Zealand. *Austral Ecology* 25: 553–562.
- Futuyma DJ 2000. Potential evolution of host range in herbivorous insects. In: van Driesche R, Heard TA, McClay AS, Reardon R eds *Host-specificity testing of exotic arthropod biological control agents: the biological basis for improvement in safety*. Morgantown, WV, USDA Forestry Service Publication FHTET-99-1. Pp. 1–10.
- Gourlay AH, Wittenberg R, Hill RL, Spiers AG, Folwer SV 2000. The biological control programme against *Clematis vitalba* in New Zealand. In: Spencer NR ed. *Proceedings of the X International Symposium on Biological Control of Weeds*, Montana State University, Bozeman, Montana, USA. Pp. 709-718.
- Greene HC 1949. Notes on Wisconsin parasitic fungi. XII. *The American Midland Naturalist* 41: 726–739.
- Greene HC 1950. Notes on Wisconsin parasitic fungi. XIV. *The American Midland Naturalist* 44: 630–642.
- Greene HC 1953. Notes on Wisconsin parasitic fungi. XVIII. *Transactions of the Wisconsin Academy of Sciences, Arts and Letters* 42: 69–81.
- Grehan JR 1989. Larval feeding habits of the Hepialidae (Lepidoptera). *Journal of Natural History* 23: 803–824
- Gressitt JL, Kimoto S 1963. The Chrysomelidae (Coleoptera) of China and Korea. Pt.2. *Pacific Insects Monograph* 1B: 743–893.
- Hamilton KGA 1997. Leafhoppers (Homoptera: Cicadellidae) of the Yukon: dispersal and endemism. In: Danks HV, Downes JA Eds *Insects of the Yukon*. Ottawa, Biological Survey of Canada (Terrestrial Arthropods). Pp. 337–375.
- Hayes L 2005. *Weed biocontrol agents for weeds in New Zealand: a field guide*. Lincoln, New Zealand, Landcare Research.
- Healy AJ 1966. Flora, adventive. In: McLintock AH ed. *An encyclopaedia of New Zealand*. . Updated 18 Sep 2007 Te Ara: the encyclopedia of New Zealand. <http://www.TeAra.govt.nz/1966/F/FloraAdventive/en>
- Hennecke B 2004. The prospect of biological control of *Mimosa pigra* with fungal pathogens in Australia In: Julien M, Flanagan G, Heard T, Hennecke B, Paynter Q, Wilson C eds *Research and management of Mimosa pigra*. Canberra, Australia, CSIRO Entomology. Pp. 117–121.
- HerbariumPDD 2008. NZFUNGI Database of New Zealand Fungi. Landcare Research, New Zealand [Web site]. Landcare Research, August 2004 2001-2008 [cited February/March 2008]. Available from <http://nzfungi.landcareresearch.co.nz>.
- Hill RJ, Foland D 1986. *Equisetum*. Poisonous plants of Pennsylvania. Harrisburg, PA, Department of Agriculture, Bureau of Plant Industry. Pp. 67–68.
- Hoffmann JH 1995. Biological control of weeds: the way forward, a South African perspective. In: Waage JK ed. *Weeds in a changing world: Proceedings of British Crop Protection Council Symposium* 64: 77–98.

- Jensen AS, Holman J 2000. *Marcrosiphum* on ferns: taxonomy, biology and evolution, including the description of three new species (Hemiptera: Aphididae). *Systematic Entomology* 25: 339–372.
- Jervis M, Vilhelmsen L 2000. Mouthpart evolution in adults of the basal, ‘symphytan’, hymenopteran lineages. *Biological Journal of the Linnean Society* 70: 121–146.
- Kirby DR, Carlson RB, Krabbenhoft KD, Mundal D, Kirby MM 2000. Biological control of leafy spurge with introduced flea beetles (*Aphthona* spp.). *Journal of Range Management* 53: 305–308.
- Kirk PM, Cannon PF, David JC, Stalpers JA eds. 2001. *Dictionary of the Fungi*. Wallingford, UK, CAB International.
- Leblanc L, Goulet H 1992. Descriptions of larvae of 8 Nearctic species of *Dolerus* (Hymenoptera: Tenthredinidae) with focus on 6 *Equisetum*-feeding species from the Ottawa region. *Canadian Entomologist* 124: 999–1014.
- Lacourt PJ 1999. Répertoire des Tenthredinidae Ouest-Palearctiques (Hymenoptera, Symphyta). *Memories de la SEF*, No: 3. Paris Société Entomologique de France.
- Large MF, Blanchon DJ Angus ML 2006. Devitalisation of imported horsetail (*Equisetum hyemale*). *New Zealand Journal of Crop and Horticultural Science* 34: 151–153.
- Liston AD 1995. *Compendium of European sawflies*. Daibersdorf 6, Gottfrieding, Germany, Chalastos Forestry.
- Liu W-C, Bai RI 2001. A new species of the genus *Cylindrosporium*. *Mycosystema* 20: 35–36.
- Louda SM, Pemberton RW, Johnson MT, Follett PA 2003. Non-target effects: the Achilles heel of biological control? Retrospective analyses to reduce risk associated with biocontrol introductions. *Annual Review of Entomology* 48: 365–396.
- Mayo GM, Roush RT 1997. Genetic variability of *Hypericum perforatum* L. (Clusiaceae) and the detection of resistance to the biological control agent *Aculus hyperici* Liro (Eriophyidae). *Plant Protection Quarterly* 12: 70–72.
- McEvoy P, Cox C, Coombs E 1991. Successful biological control of ragwort, *Senecio jacobaea*, by introduced insects in Oregon. *Ecological Applications* 1: 430–442.
- McFadyen REC 1998. Biological control of weeds. *Annual Review of Entomology* 43: 369–393.
- Medvedev LN, Roginskaja EJ 1988. *Catalogue of host plants of leaf-beetles of USSR*. Moscow: 192 pp. {In Russian}
- Mel'nik VA 2000. *Key to the fungi of the genus Ascochyta* Lib. (Coelomycetes). 2nd ed. Berlin, Biologischen Bundesanstalt für Land- und Forstwirtschaft.
- Memmott J, Fowler SV, Hill RL 1998. The effect of release size on the probability of establishment of biological control agents: gorse thrips (*Sericothrips staphylinus*) released against gorse (*Ulex europaeus*) in New Zealand. *Biocontrol Science and Technology* 8: 103–115.
- Memmott J, Craze PG, Harman HM, Syrett P, Fowler SV 2005. The effect of propagule size on the invasion of an alien insect. *Journal of Animal Ecology* 74: 50–62.
- Morin L, Edwards PB 2006. Selection of biological control agents for bridal creeper: a retrospective review. *Australian Journal of Entomology* 45: 287–291.
- Morris MG 2002. True weevils (Part 1) Coleoptera: Curculionidae (subfamilies Raymondionyminae to Smicronychinae). *Handbooks for the identification of British insects* 5 (17b). Pp. 1–149.
- Nickel H, Remane R 2002. *Artenliste der Zikaden Deutschlands, mit Angabe von Nährpflanzen, Nahrungsbreite, Lebenszyklus, Areal und Gefährdung* (Hemiptera, Fulgoromorpha et Cicadomorpha). *Beiträge zur Zikadenkunde* 5: 27–64. Translation available as pdf at (<http://www.gwdg.de/~hnickel>).
- Page AR, Lacey KL 2006. *Economic Impact Assessment of Australian Weed Biological Control*. (Technical Series #10). Cooperative Research Centre for Australian Weed Management, Adelaide, Australia.

- Paynter Q, Flanagan GJ 2004. Integrating herbicide and mechanical control treatments with fire and biological control to manage an invasive wetland shrub, *Mimosa pigra*. *Journal of Applied Ecology* 41: 615–629.
- Paynter QE, Fowler SV, Gourlay AH, Haines ML, Harman HM, Hona SR, Peterson PG, Smith LA, Wilson-Davey JRA, Winks CJ, Withers TM 2004. Safety in New Zealand weed biocontrol: a nationwide survey for impacts on non-target plants. *New Zealand Plant Protection* 57: 102–107.
- Paynter Q, Waipara N, Peterson P, Hona S, Fowler S, Gianotti A, Wilkie P 2006. The impact of two introduced biocontrol agents, *Phytomyza vitalbae* and *Phoma clematidina*, on *Clematis vitalba* in New Zealand. *Biological Control* 36: 350–357.
- Paynter Q, Martin N, Berry J, Hona S, Peterson P, Gourlay AH, Wilson-Davey J, Smith L, Winks C, Fowler SV 2008. Non-target impacts of *Phytomyza vitalbae* a biological control agent of the European weed *Clematis vitalba* in New Zealand. *Biological Control* 44: 248–258.
- Pemberton RW 2000. Predictable risk to native plants in weed biological control. *Oecologia* 125: 489–494.
- Pryer KM, Schuettpelez E, Wolf PG, Schneider H, Smith AR, Cranfill R 2004. Phylogeny and evolution of ferns (monilophytes) with a focus on the early leptosporangiate divergences. *American Journal of Botany* 91: 1582–1598.
- Punithalingam E 1988. Ascochyta II. Species on monocotyledons (excluding grasses), cryptogams and gymnosperms. *Mycological Papers* 159: 1–235.
- Radulovic N, Stojanovic G, Palic R. 2005. Composition and antimicrobial activity of *Equisetum arvense* L. essential oil. *Phytotherapy Research* 20(1): 85–88.
- Rai MK 1990. New records of fungi from India. *Indian Journal of Mycology & Plant Pathology* 20: 199–201.
- Robbins R 1990. Provisional atlas of the leaf miners of Warwickshire, A, with notes on others occurring in the midlands. Warwick UK, Warwickshire Museum Service.
- Rosenthal SS 1996. Chapter 4.1.1 *Aceria*, *Epitrimerus* and *Aculus* species and biological control of weeds. In: Lindquist EE, Sabelis MW, Bruin J eds *World crop pests*, Vol. 6. Eriophyoid mites: their biology, natural enemies and control. Amsterdam, Elsevier Science BV. Pp. 729–739.
- Roy B, Popay I, Champion P, James T, Rahman A 1998. An illustrated guide to common weeds of New Zealand. Meredith, Australian Print Group, New Zealand Plant Protection Society.
- Schuettpelez E, Korall P, Pryer KM 2006. Plastid atpA data provide improved support for deep relationships among ferns. *Taxon* 55: 897–906.
- Shin HD 1994. New fungal diseases of economic resource plants in Korea (1). *The Plant Pathology Journal* 9: 181–191.
- Shoemaker RA, Babcock CE 1989. Phaeosphaeria. *Canadian Journal of Botany* 67: 1500–1599.
- Söderman G 2007. Taxonomy, distribution, biology and conservation status of Finnish Auchenorrhyncha (Hemiptera: Fulgoromorpha et Cicadomorpha). *The Finnish Environment* 7. Helsinki, Finnish Environment Institute.
- Spencer KA 1972. Diptera, Agromyzidae. *Handbooks for the identification of British Insects*, vol X part 5(g), Royal Entomological Society of London.
- Stewart CA, Chapman RB, Frampton CMA 2000. Growth of alligator weed (*Alternanthera philoxeroides* (Mart.) Griseb. (Amaranthaceae)) and population development of *Agasicles hygrophila* Selman & Vogt (Coleoptera: Chrysomelidae) in northern New Zealand. *Plant Protection Quarterly* 15: 95–101.
- Syrett P, Fowler SV, Emberson RM 1996. Are chrysomelid beetles effective agents for biological control of weeds? In: Moran VC, Hoffmann JH eds *Proceedings of the IX International Symposium on Biocontrol of Weeds*, Stellenbosch, South Africa, University of Cape Town. Pp. 399–407.

- Taeger A, Blank SM 1998. Pflanzenwespen Deutschlands (Hymenoptera, Symphyta). Keltorn, Goecke und Evers.
- Thum R, Paynter Q, Völkl W, Hoffmann KH 1997. Ant-Homoptera-Phytophage-plant interaction: are ant-tended Homoptera good for biological control? *Mitteilung Deutschen Gesellschaft für allgemeine und angewandte Entomologie* 11: 711–715.
- van Klinken RD, Heard TA 2000. Estimating fundamental host range: a host-specificity study of a potential biocontrol agent for *Prosopis* species (Leguminosae). *Biocontrol Science and Technology* 10: 331–342.
- van Klinken RD, Fichera G, Cordo H, 2003. Targeting biological control across diverse landscapes: the release, establishment, and early success of two insects on mesquite (*Prosopis* spp.) insects in Australian rangelands. *Biological Control* 26: 8–20.
- Wapshere AJ 1974. A strategy for evaluating the safety of organisms for biological weed control. *Annals of Applied Biology* 77: 201–211.
- Webb CJ, Sykes WR, Garnock-Jones, PJ, 1988. *Flora of New Zealand Volume 4: naturalised Pteridophytes, Gymnosperms, Dicotyledons*. Christchurch, Botany Division, Department of Scientific and Industrial Research.
- Webb CJ, Sykes WR, Garnock-Jones PJ, Brownsey PJ 1995. Checklist of dicotyledons, gymnosperms, and pteridophytes naturalised or casual in New Zealand: additional records 1988-1993. *New Zealand Journal of Botany* 33: 151–182.
- Weiffenbach H 1985. Symphyta (Hymenoptera) von Süd-Niedersachsen, Nord- und Mittelhessen. *Mitteilungen der Münchener Entomologischen Gesellschaft* 75: 5–44.
- Wilson SW, Mitter C, Denno RF, Wilson MR 1994. Evolutionary patterns of host plant use by delphacid planthoppers and their relatives. In: Denno RF, Perfect TJ eds *Planthoppers: their ecology and management*. New York, Chapman and Hall. Pp. 7–113.
- Wikstrom N, Pryer KM 2005. Incongruence between primary sequence data and the distribution of a mitochondrial *atp1* group II intron among ferns and horsetails. *Molecular Phylogenetics and Evolution* 36: 484–493.

Appendices

Appendix 1: Fungi recorded on field horsetail *Equisetum arvense* and their potential usefulness for biological control

Phylum, Order, Family	Species (and other names) ¹	Range on field horsetail ²	Likely to be damaging?	Likely to be host-specific? (and comments)	Found in NZ/biocontrol potential?
Ascomycota Dothideales Dothideaceae	<i>Scirrhia castagnei</i>	France; Scotland (Mid Ebudes)	Unknown: Reference is a list of fungi in the Hebrides. No mention of disease symptoms (Dennis 1986)	Yes: 6 records in FRDBI ³ . All on 3 <i>Equisetum</i> species (incl. <i>E. arvense</i>)	No/?
Ascomycota Helotiales Helotiaceae	<i>Hymenoseyphus ochroleucus</i>	England (Surrey)	Unknown	Unknown: Only 1 record in British fungi: On/with <i>E. arvense</i>	No/?
Ascomycota Helotiales Helotiaceae	<i>Hymenoseyphus rhodoleucus</i>	England (Gloucestershire, Lancaster, Northumberland, Walkers Heath, Worcestershire); Scotland (Mid Ebudes, Wigtownshire); Wales (Pembrokeshire)	No: Saprobe ⁴ found on/with litter, dead material, dead stems and stems	Probably: 39 records in FRDBI, 37 of them on <i>Equisetum</i> species, 1 host not stated, and 1 on dead stems of <i>Calluna vulgaris</i> (Heather)	No/No
Ascomycota Helotiales Helotiaceae	<i>Stannaria personii</i> (also recorded as <i>S. equiseti</i> {old name}) anamorph = <i>Titaospora equiseti</i> (anamorph also recorded as <i>Gloeosporium equiseti</i> , and <i>Cylindrosporium</i>	Austria; British Isles; Canada (Ontario); Czech Republic; France; Germany; Italy; Korea; USA (Idaho, Washington, South Dakota). Also found widely in the northern hemisphere on other	Possibly: Recorded: on/with host, on stems and leaves, on living leaves, on withered stems, on dead stems & on stem spots. Reported to cause leaf spot and leaf blight on <i>E. arvense</i> in	Probably: 96 records, 18 host/pathogen combinations in FDSM ⁵ and 16 records in FRDBI. All except 2 collections on <i>Equisetum</i> spp. Exceptions are on <i>Ilex cornuta</i> (Dicotyledon, same genus as Holly) and <i>Lagerstroemia</i>	No, apart from collections in PDD (see main text)/Yes

Phylum, Order Family	Species (and other names) ¹	Range on field horsetail ²	Likely to be damaging?	Likely to be host-specific? (and comments)	Found in NZ/biocontrol potential?
	<i>equiseti</i> {old names}}	<i>Equisetum</i> spp.	Korea (Shin 1994)	<i>indicia</i> (Crêpe myrtle) in Florida	
Ascomycota Helotiales Helotiaceae	<i>Stannaria</i> sp. anamorph = <i>Titaospora</i> sp.	USA (California)	Unknown	Unknown: Just 1 record in FDSM	?/?
Ascomycota Helotiales Helotiaceae	<i>Titaospora detospora</i> teleomorph = <i>Stannaria</i> sp.	USA (Wisconsin, Wyoming); Poland; West Germany; Czechoslovakia	Unknown. Recorded as on stems or leaves, reference does not state if pathogenic (Greene 1949)	Yes: 9 records in FDSM, all on 5 <i>Equisetum</i> spp. (incl. <i>E. arvense</i>). Lives in temperate climates (Greene 1949)	No/?
Ascomycota Helotiales Hyaloscyphaceae	<i>Lachnella alboviolascens</i> (Note that <i>Lachnella</i> is said to be a synonym of <i>Lachnum</i> in Kirk et al. 2001)	Canada	Probably not: Saprobe?	No: Many distantly related hosts. Widespread on woody and herbaceous stems	Yes, indigenous/No
Ascomycota Helotiales Hyaloscyphaceae	<i>Lachnella</i> (= <i>Lachnum</i> ?) <i>villosa</i>	Wales (Brecon)	No: Saprobe, where host given, mostly on dead material	No: 273 records in FRDBI, hosts mostly not given	Yes, exotic/No
Ascomycota Helotiales Hyaloscyphaceae	<i>Lachnum virgineum</i>	Scotland (Wigtownshire)	No: Saprobe, where host given, on dead material	No: 2276 records in FRDBI, hosts mostly not given	Yes/No
Ascomycota Helotiales Hyaloscyphaceae	<i>Psilachnum inquilinum</i>	England (Hertfordshire, Northumberland, Surrey, Yorkshire); Scotland (Wigtownshire); Wales (Cardiganshire, Pembrokeshire)	No: recorded on/with host, stems, dead stems & dead material. Probably saprobe	No: Mostly found on dead tissues of 5 <i>Equisetum</i> species. However, also less commonly found on dead material of unrelated plants such as <i>Populus tremula</i> (aspens) and <i>Abies</i> (fir)	No/No
Ascomycota Hypocreales	<i>Lasionectria sylvana</i> (also recorded as <i>Nectria</i>)	USA (New York)	Unknown	No: 14 records in FDSM, 9 of these from NZ. Hosts include	Yes/No

Phylum, Order Family	Species (and other names) ¹	Range on field horsetail ²	Likely to be damaging?	Likely to be host-specific? (and comments)	Found in NZ/biocontrol potential?
Bionectriaceae	<i>sylvana</i> {old name}}			ferns and monocots (e.g., <i>Astelia</i> , <i>Rhopalosiphum sapida</i> {nikau} & <i>Phormium tenax</i> {flax})	
Ascomycota Hypocreales Incertae sedis ⁶	<i>Verticillium dahliae</i>	Canada	Yes: Causes wilt disease in many plants (Kirk et al. 2001)	No: 51 records in FRDBI, 343 in FDSM. Found on hosts in many different families, cosmopolitan	Yes, exotic/ No
Ascomycota Hypocreales Nectriaceae	<i>Acremonium boreale</i> teleomorph = <i>Nectria tuberculariformis</i>	Canada	Yes. Listed on MAFs unwanted organisms register as 'regulated pest' (HerbariumPDD 2001–2008)	No: Many unrelated hosts	No/ No
Ascomycota Incertae sedis	<i>Ascochyta</i> cf. <i>teretiuscula</i>	England (Surrey)	No: Found on dead stem so may be saprobe.	?: No more information because species ID uncertain	No/?
Ascomycota Incertae sedis	<i>Ascochyta equiseti</i> (also recorded as <i>Stagonosporopsis equiseti</i> {old name})	Belgium; Czechoslovakia; Denmark; England (Lancaster, London, Warwickshire, Yorkshire); France; Germany; Italy; Romania; Switzerland; USA (Wisconsin) & the USSR China, Poland, Germany	Yes: Said to be definitely parasitic (Greene 1950). Associated with dead and dying leaves and stems (Mel'nik 2000) or with indistinct lesions on bleached areas on stems (Punithalingam 1988)	Yes: 21 records in FRDBI, 19 records FDSM, all from various <i>Equisetum</i> species. (Mel'nik 2000; Punithalingam 1988)	No/ Yes
Ascomycota Incertae sedis	<i>Cylindrosporium equiseti</i>		Maybe: Recorded in living foliage, associated with small spots 0.5–8 mm in diameter (Lui and Bai 2001)	Yes: 5 collections, 4 from <i>E. arvense</i> , 1 from <i>E. sp.</i> Probably genus specific. This species (<i>C. equiseti</i> W.C. Liu et R.L. Bai) was described recently and is the only <i>Cylindrosporium</i>	No/ Yes

Phylum, Order Family	Species (and other names) ¹	Range on field horsetail ²	Likely to be damaging?	Likely to be host-specific? (and comments)	Found in NZ/biocontrol potential?
Ascomycota Incertae sedis	<i>Gyrothrix verticillata</i>	England (Surrey)	Unknown	known from the Equisetaceae (Lui & Bai 2001). An earlier fungus initially called <i>Cylindrosporium equiseti</i> (Desm.) Died is a synonym of <i>Titaeospora equiseti</i> (Desm.) Vassiljevsky (teleomorph = <i>Stannaria persoonii</i>)	
Ascomycota Incertae sedis	<i>Monodidymaria equiseti</i> (also recorded as <i>Cercospora equiseti</i> {old name})	Russia; USSR	Probably: "Segments or entire branchlets of the stems discoloured, pale to brown" (Braun 1994)	No: Other hosts unrelated (e.g., <i>Fagus sylvatica</i> (Beech), <i>Phormium tenax</i> (NZ Flax)) Possibly: only 2 records in FDSM, both on <i>E. arvense</i> . Reference describes fungus as 'insufficiently known' (Braun 1994)	No/No No/Yes
Ascomycota Incertae sedis	<i>Septogloeum equiseti</i>	England (Norfolk)	Probably not: listed on dead stems so probably saprobe	Probably: 2 records in FRDBI, one on <i>E. arvense</i> and one on <i>E. palustre</i>	No/?
Ascomycota Incertae sedis	<i>Stagonospora</i> sp.	USA (Wisconsin)	Unknown	Unknown: Just 1 record in FDSM. No more information as because species ID uncertain	?/?
Ascomycota Microascales Ceratocystidaceae	<i>Thielaviopsis basicola</i>	Canada	Yes: Causes black root rot (Kirk et al. 2001)	No: Widespread, cosmopolitan, hosts in many plant families	Yes, exotic/No
Ascomycota Mycosphaerellales Mycosphaerellaceae	<i>Cladosporium cladosporioides</i>	England (Surrey)	No: Saprobe	No: Many unrelated plant hosts & organic materials such as dung, mattresses, walls	Yes, origin/No uncertain
Ascomycota	<i>Cladosporium herbarum</i>	USA (Iowa)	No: Saprobe	No: Very common	Yes, exotic/No

Phylum, Order, Family	Species (and other names) ¹	Range on field horsetail ²	Likely to be damaging?	Likely to be host-specific? (and comments)	Found in NZ/biocontrol potential?
Mycosphaerellales Mycosphaerellaceae	teleomorph = <i>Davidiella tassiana</i>			cosmopolitan sp., hundreds of known hosts (mostly on dead tissues), and frequently isolated from air, soil, foodstuffs etc. (Ellis 1971)	
Ascomycota Mycosphaerellales Mycosphaerellaceae	<i>Heterosporium equiseti</i> (Note that <i>Heterosporium</i> is said to be a synonym of <i>Cladosporium</i> in Kirk et al. 2001)	USA (Wisconsin)	Probably: Described on brown distal portions of branchlets “fungus develops progressively inward from branchlet tips until the entire plant may be involved” Greene 1950)	Yes: 4 records in FDSM: 2 from <i>E. arvense</i> , 2 from two difference subspecies of <i>E. sylvaticum</i> . Probably genus specific	No/Yes
Ascomycota Mycosphaerellales Mycosphaerellaceae	<i>Mycosphaerella altera</i>	USA (Wisconsin)	Probably: Recorded on dead tips of living branches (Greene 1953)	Probably: 4 records in FDSM: 1 on <i>E. arvense</i> ; 2 on <i>E. hymalea</i> ; 1 on <i>E. telmateia</i>	No/Yes
Ascomycota Mycosphaerellales Mycosphaerellaceae	<i>Mycosphaerella aspidii</i>	Sweden	Unknown (References = taxonomic or geographic lists: not helpful)	Probably specific to ferns and fern allies: 8 records in FRDBI: 4 on <i>Equisetum</i> species including <i>E. arvense</i> : 3 on true ferns (<i>Dryopteris</i> sp.; <i>Pteridium</i> spp. & <i>Nephrodium filix-mas</i>)	No/?
Ascomycota Mycosphaerellales Mycosphaerellaceae	<i>Mycosphaerella equiseticola</i>	Sweden	Unknown (References = geographic lists: not helpful)	Yes: 8 records in FDSM, all on <i>Equisetum</i> species (7 different <i>E.</i> spp. including <i>E. arvense</i>)	No/?
Ascomycota Pleosporales Diademaceae	<i>Clathrospora diplospora</i>	Poland	Unknown	No: more than 100 known hosts	No/No
Ascomycota	<i>Stagonospora equiseti</i>	England (Yorkshire)	Probably not: listed on	Probably: 5 records in FRDBI,	No/?

Phylum, Order, Family	Species (and other names) ¹	Range on field horsetail ²	Likely to be damaging?	Likely to be host-specific? (and comments)	Found in NZ/biocontrol potential?
Pleoporales Incertae sedis			dead stems in 2 of 5 records so probably saprobe	all on <i>Equisetum</i> spp.	
Ascomycota Pleoporales Leptosphaeriaceae	<i>Leptosphaeria arvensis</i> (also listed under name of anamorph <i>Heptameria arvensis</i>)	USA (California); Italy	Unknown: recorded on stems and/or dead stems, no more info. available in literature.	Yes: 5 records FDSM: 4 from <i>E. arvensis</i> , 1 from <i>E. hyemale</i>	No/?
Ascomycota Pleoporales Phaeosphaeriaceae	<i>Phaeosphaeria equiseti</i>	Norway	Unknown. Reference does not say if pathogenic (Shoemaker and Babcock 1989)	12 records in FDSM, all on <i>Equisetum</i> sp. (4 difference <i>E.</i> spp. including <i>E. arvensis</i>)	No/?
Ascomycota Pleoporales Phaeosphaeriaceae	<i>Phaeosphaeria fuckelii</i>	Sweden	Unknown. Reference does not say if pathogenic (Shoemaker and Babcock 1989)	No: more than 40 hosts. Mostly grasses, but also other monocots, ferns and fern allies. ITS sequence puts this fungus with other <i>Phaeosphaeria</i> found on these types of hosts (Camara et al. 2001)	Yes, exotic. Specimens on two grasses in PDD/No
Ascomycota Pleoporales Phaeosphaeriaceae	<i>Phaeosphaeria lindii</i>	Norway	Unknown. Reference does not say if pathogenic (Shoemaker and Babcock 1989)	Yes: 6 records in FDSM, all on <i>Equisetum</i> sp. (4 difference <i>E.</i> spp. including <i>E. arvensis</i>)	No/?
Basidiomycota Agaricales Flagelloscyphaceae	<i>Flagelloscypha minutissima</i>	England (Surrey)	No: Saprobe	No: more than 10 distantly related hosts & wood, bark etc.	No/No
Basidiomycota Agaricales Pluteaceae	<i>Volvariella gloiocephala</i>	England (Gloucestershire)	No: Mushroom, saprobe	No: 961 records in British fungi, substrate mostly not given, but where given = soil, sand, on/with grasses	No/No

Phylum, Order, Family	Species (and other names) ¹	Range on field horsetail ²	Likely to be damaging?	Likely to be host-specific? (and comments)	Found in NZ/biocontrol potential?
Basidiomycota Ceratobasidiales Ceratobasidiaceae	<i>Rhizoctonia solani</i> teleomorph = <i>Thanatephorus cucumeris</i>	UA (Illinois, Minnesota, Texas)	Yes: Causes root rot. Listed by MAF as a 'regulated pest' (HerbariumPDD 2001-2008)	No: More than 2400 fungus/host combinations. Cosmopolitan, plurivorous, commonly found on roots but also stems, seedlings, leaves and fruits	Yes, exotic/ No
Zygomycota Mucorales Choanephoraceae	<i>Poitrasia circinans</i>	India	Probably not: Described on roots of living plant (Rai 1990), and this seems to be the only record	Unknown: Only the single record on <i>E. arvense</i> in FDSM	No/?

¹ Many fungi have more than one Latin name because they can produce more than one type of spore. The name given when they are producing 'sexual' spores is called the teleomorph, whereas the stage producing 'asexual' spores is called the anamorph. The two stages often look completely different. Fungi are classified according to their 'teleomorph' name, unless the 'anamorph' is the only form known. So, appendix 1 gives the taxonomy of the teleomorph, but column 2 uses whichever name/names were recorded when the fungus was found on *E. arvense*. If a fungus was listed under an out-of-date name (synonym) this is also stated in column 2.

² Only the places where the organism was found associated with *E. arvense* are listed here. It may also be found elsewhere on other hosts.

³ FRDBI = the Fungal Records Database of Britain and Ireland (FRDBI) at <http://194.203.77.76/fieldmycology/FRDBI/FRDBI.asp>

⁴ Saprobe: An organism using dead organic material as food and commonly causing its decay (Kirk *et al.* 2001). Unlikely to cause disease and therefore probably insufficiently damaging to be useful for biocontrol.

⁵ FDSM = USDA Fungus-host database at <http://nt.ars-grin.gov/fungal/databases/fungushost/FungusHost.cfm>

⁶ Insertae sedis = of uncertain taxonomic position within a higher taxonomic order (e.g., Phylum known, but order within that phylum uncertain).

Appendix 2: Records of invertebrates feeding on field horsetail *Equisetum arvense*

Order and Family	Species	Type of organism	Range	Likely to be sufficiently host-specific
TYLENCHIDA				
Pratylenchidae	<i>Pratylenchus penetrans</i>	Nematode worm	Cosmopolitan (including NZ)	No: Extremely polyphagous nematode crop pest
PROSTIGMATA				
Eriophyidae	<i>Eriophyes equiseti</i>	Eriophyid mite	Hungary	Yes: Only recorded from <i>E. arvense</i> (Farkas 1960)
Tetranychidae	<i>Bryobia lagodechiana</i>	Spider mite	Cosmopolitan (including NZ)	No: Polyphagous (http://www.montpellier.inra.fr/CBGP/spmweb/advanced.php)
Tetranychidae	<i>Tetranychus urticae</i>	Spider mite	Cosmopolitan (including NZ)	No: Extremely polyphagous (<i>E. arvense</i> host record: Aucejo et al. 2003; cited in Spider Mites Web database http://www.montpellier.inra.fr/CBGP/spmweb/advanced.php)
HEMIPTERA				
Pseudococcidae	<i>Maconelliococcus hirsutus</i>	Mealybug	Cosmopolitan (but not NZ)	No: Extremely polyphagous pest (Pink hibiscus mealybug)
Aphididae	<i>Macrosiphum</i> = <i>Sitobion equiseti</i>	Aphid	Holarctic	Yes: Specific to <i>Equisetum</i> (Jensen & Holman 2000)
Cicadellidae	<i>Macrosteles borealis</i> <i>Macrosteles frontalis</i> <i>Notus flavipennis</i>	Leafhopper Leafhopper Leafhopper	Nearctic Europe Palearctic	Yes: Host plants <i>Equisetum</i> spp. (Hamilton 1997) Yes: Host plants <i>E. arvense</i> , <i>E. palustre</i> (Nickel & Remane 2002) No: Host plants <i>Equisetum</i> , <i>Carex versicaria</i> , <i>Scirpus sylvaticus</i> (Söderman 2007)
	<i>Ophiolix paludosa</i>	Leafhopper	Europe	No: Host plants <i>Carex</i> and <i>Equisetum</i> (Söderman 2007)
Delphacidae	<i>Javesella kilmani</i> <i>Javesella stali</i>	Planthopper Planthopper	Canada; USA Palearctic	Yes: Host plant <i>Equisetum</i> sp. (Wilson et al. 1994) Yes: Monophagous on <i>E. arvense</i> (Nickel & Remane 2002)

HYMENOPTERA

Tenthredinidae

<i>Ametastegia equiseti</i>	Sawfly	Palaearctic	No: Polyphagous (host plants chiefly Polygonaceae; Weiffenbach 1985) Yes: Host plant <i>E. arvensis</i> (Leblanc & Goulet 1992)
<i>Dolerus acidus</i>	Sawfly	Canada (Ottawa region) Europe	Yes: Host plant <i>Equisetum</i> including <i>E. arvensis</i> (Weiffenbach 1985); very common in England (Benson 1952)
<i>Dolerus aericeps</i>	Sawfly	Canada (Ottawa region)	Yes: Host plant <i>E. arvensis</i> (Leblanc & Goulet 1992)
<i>Dolerus apricus</i>	Sawfly	Canada (Ottawa region)	Yes: Host plant <i>E. arvensis</i> (Leblanc & Goulet 1992)
<i>Dolerus aprilis</i>	Sawfly	Canada (Ottawa region) Palaearctic	Yes: Host plant <i>Equisetum</i> including <i>E. arvensis</i> (Benson 1952; Weiffenbach 1985)
<i>Dolerus bimaculatus</i>	Sawfly	Palaearctic	Yes: Host plant <i>E. palustre</i> , <i>E. arvensis</i> (Liston, 1995; Lacourt 1999)
<i>Dolerus eversmanni</i>	Sawfly	Holarctic	Yes: Host plant <i>E. palustre</i> , <i>E. arvensis</i> (Weiffenbach 1985);
<i>Dolerus germanicus</i>	Sawfly	Palaearctic	Çalmaşur & Özbek 2004)
<i>Dolerus gessneri</i>	Sawfly	Northern Palaearctic	Yes: Host plant <i>Equisetum</i> (Benson 1952)
<i>Dolerus gilvipes</i>	Sawfly	Holarctic	Yes: Host plants <i>E. pratense</i> and <i>E. sylvaticum</i> (Weiffenbach 1985) and <i>E. arvensis</i> (subspecies <i>albifrons</i> ; Leblanc & Goulet 1992)
<i>Dolerus pratensis</i>	Sawfly	Holarctic	Yes: Host plant <i>Equisetum</i> spp., <i>E. heliocharis</i> (Liston, 1995; Lacourt, 1999); often abundant in Britain & Ireland (Benson 1952)
<i>Dolerus pratorum</i>	Sawfly	Palaearctic	Yes: Host plant <i>Equisetum</i> , including <i>E. arvensis</i> (Benson 1952; Weiffenbach 1985)
<i>Dolerus subfasciatus</i>	Sawfly	Canada (Ottawa region)	Yes: Host plant <i>E. arvensis</i> (Leblanc & Goulet 1992)
<i>Dolerus tibialis conjugatus</i>	Sawfly	Canada (Ottawa region)	Yes: Host plant <i>E. arvensis</i> (Leblanc & Goulet 1992)
<i>Dolerus vestigialis</i>	Sawfly	Holarctic	Yes: Host plants <i>E. palustre</i> , <i>E. sylvaticum</i> , <i>E. arvensis</i> , <i>E. pratense</i> (Liston, 1995; Taeger & Blank, 1998; Lacourt, 1999).
<i>Dolerus yukonensis</i>	Sawfly	Holarctic	Yes: Host plant <i>Equisetum</i> (Benson 1952)

COLEOPTERA									
Chrysomelidae	<i>Hippuriphila modeeri</i>	Flea beetle	W Palaearctic to Mongolia Japan, Korea, N. China	Yes: Host plants <i>E. arvense</i> (Medvedev & Roginskaja 1988); <i>Equisetum</i> spp. <i>E. palustre</i> (Bullock 1992). No: Host plants <i>E. arvense</i> , <i>Hosta</i> sp. (Agavaceae) (Gressitt & Kimoto 1963).					
	<i>Liprus punctatostrigatus</i>	Flea beetle							
Erihniidae	<i>Grypus equiseti</i>	Weevil	Holarctic; widespread in Britain (Morris 2002)	Yes: Host plants <i>E. palustre</i> and <i>E. arvense</i> (Morris 2002)					
DIPTERA									
Agromyzidae	<i>Liriomyza occipitalis</i>	Leaf-mining fly	Europe	Yes: Host plant <i>E. arvense</i> (Spencer 1972)					
	<i>Liriomyza equiseti</i>	Leaf-mining fly	Holarctic	Yes: Host plant <i>E. arvense</i> in stems and thicker branches; larva eats some pith as well as parenchyma (Spencer 1972; Robbins 1990)					
	<i>Liriomyza virgo</i>	Leaf-mining fly	Holarctic	Yes: Main host plant <i>E. fluviatile</i> , also recorded from <i>E. arvense</i>					
LEPIDOPTERA									
Noctuidae	<i>Hydraecia micacea</i>	Moth	Native to Europe, introduced North America	No: Highly polyphagous (NHM HOSTS database)					
	<i>Xylota vetusta</i>	Moth	Europe	No: Highly polyphagous (NHM HOSTS database)					
Hepialidae	<i>Endoclita excrescens</i>	Moth	East Asia	No: Highly polyphagous tunnels branches (Grehan 1989)					
	<i>Triodia sylvina</i>	Moth	Europe	No: Highly polyphagous root-feeder					
Tortricidae	<i>Loxoterma tiedemanniana</i>	Moth	Palaearctic (Finland, South Siberian mountains)	Yes: Host plant <i>Equisetum</i> (in stems; NHM HOSTS database)					

Appendix 3: Steps in a Biocontrol Project

A classical biocontrol programme typically works through the following steps; this is usually done in a sequential manner, but some activities may occur concurrently.

1. Explore the feasibility of project. If project looks feasible, proceed.
2. Survey weed in places where biocontrol is desired. If any potential agents are found explore ways to maximise them. If any likely impediments are found look for ways to mitigate them.
3. Undertake molecular studies of the weed to help narrow down the best place in the native range to find natural enemies.
4. Unless natural enemies are already well known, survey weed in native range. Identify and study life cycles of natural enemies found.
5. Determine host range for potential agents. Abandon any species that do not appear to be safe or effective enough.
6. Apply to authorities for permission to release agents.
7. If permission is granted import, clear through quarantine, and develop rearing techniques for new agents (if not already known)
8. Mass rear and release agents over several years.
9. Monitor establishment success and dispersal of agents over several years.
10. Harvest and redistribute agents
11. Evaluate success of project. Decide if further agents are needed.

An inundative biocontrol programme where mycoherbicides are developed for use, typically works through the following steps in a sequential manner:

1. Survey the weed to look for potentially useful pathogens (if not already known)
2. Undertake glasshouse trials to explore host range and pathogenicity
3. Develop prototype formulation and test under field conditions
4. Refine formulation
5. Explore techniques for mass production and prospects for commercialisation
6. Undertake risk analysis and apply for registration